



Luminous dark photon clouds



Cristina Mondino



Gravitational Wave Probes of Black Hole Environments

IFPU focus week, 26th of May 2025

Phys. Rev. D 107 (2023) 7, arxiv:2212.09772

Phys. Rev. D 111 (2025) 8, arxiv:2501.02052

With: Nils Siemonsen, Lorenzo Mirasola, Daniel Egaña-Ugrinovic, Junwu Huang, Masha Baryakhtar, William E.

East + CW group at INFN Roma/La Sapienza



Model and motivations

Standard Model of Elementary Particles

three generations of matter (fermions)						interactions / force carriers (bosons)	
I			II			III	
mass charge spin	$\approx 2.16 \text{ MeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$	$\approx 1.273 \text{ GeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$	$\approx 172.57 \text{ GeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$	0 0 1	$\approx 125.2 \text{ GeV}/c^2$ 0 0		
	u up	c charm	t top	g gluon	H higgs		
QUARKS	$\approx 4.7 \text{ MeV}/c^2$ $-\frac{2}{3}$ $\frac{1}{2}$	$\approx 93.5 \text{ MeV}/c^2$ $-\frac{2}{3}$ $\frac{1}{2}$	$\approx 4.183 \text{ GeV}/c^2$ $-\frac{2}{3}$ $\frac{1}{2}$	0 0 1			
	d down	s strange	b bottom	γ photon			
LEPTONS	$\approx 0.511 \text{ MeV}/c^2$ -1 $\frac{1}{2}$	$\approx 105.66 \text{ MeV}/c^2$ -1 $\frac{1}{2}$	$\approx 1.77693 \text{ GeV}/c^2$ -1 $\frac{1}{2}$	$\approx 91.188 \text{ GeV}/c^2$ 0 1			
	e electron	μ muon	τ tau	Z Z boson			
	$< 0.8 \text{ eV}/c^2$ 0 $\frac{1}{2}$	$< 0.17 \text{ MeV}/c^2$ 0 $\frac{1}{2}$	$< 18.2 \text{ MeV}/c^2$ 0 $\frac{1}{2}$	$\approx 80.3692 \text{ GeV}/c^2$ ± 1 1			
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson			

SCALAR BOSONS

GAUGE BOSONS
VECTOR BOSONS



$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^\mu A'_\mu - \frac{\epsilon}{2}F^{\mu\nu}F'_{\mu\nu}$$

- New U(1) with massive vector boson gauge field
- Can mix kinetically with the SM photon
- Can be light
- Can be the DM or SM-DM mediator

Model and motivations

Standard Model of Elementary Particles



	three generations of matter (fermions)			interactions / force carriers (bosons)	
	I	II	III		
mass	$\approx 2.16 \text{ MeV}/c^2$	$\approx 1.273 \text{ GeV}/c^2$	$\approx 172.57 \text{ GeV}/c^2$	0	$\approx 125.2 \text{ GeV}/c^2$
charge	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0	0
spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0
	u up	c charm	t top	g gluon	H higgs
	d down	s strange	b bottom	γ photon	
	e electron	μ muon	τ tau	Z Z boson	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	

QUARKS (purple box)

LEPTONS (green box)

SCALAR BOSONS (yellow box)


GAUGE BOSONS VECTOR BOSONS (red box)



$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^{\mu}A'_{\mu} + J_{\text{EM}}^{\mu}(A_{\mu} + \boxed{\varepsilon}A'_{\mu})$$

- New U(1) with massive vector boson gauge field
- Can mix kinetically with the SM photon
- Can be light
- Can be the DM or SM-DM mediator


Dark photon parameter space



From Ciaran O'Hare (cajohare.github.io/AxionLimits/)

Dark photon parameter space

If
 $\Omega_{A'} \ll \Omega_{\text{DM}}$



From Ciaran O'Hare (cajohare.github.io/AxionLimits/)

Black hole superradiance - vector field


M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019

...

New massive vector boson

$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^\mu A'_\mu$$

$$\frac{h}{mc} \sim \frac{G_N M}{c^2} \quad \sim$$



Black hole superradiance - vector field

M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019


...

New massive vector boson

$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^{\mu}A'_{\mu}$$

$$\frac{h}{mc} \sim \frac{G_N M}{c^2} \quad \sim$$

- BH loses energy and angular momentum in favor of amplification of boson wave



Black hole superradiance - vector field

M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019


...

New massive vector boson

$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^\mu A'_\mu$$

$$\frac{h}{mc} \sim \frac{G_N M}{c^2} \quad \sim$$

- BH loses energy and angular momentum in favor of amplification of boson wave
- Create large bosonic cloud around spinning BHs



Black hole superradiance - vector field


M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019

...

New massive vector boson

$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^{\mu}A'_{\mu}$$

$$\frac{h}{mc} \sim \frac{G_N M}{c^2} \quad \sim$$



- BH loses energy and angular momentum in favor of amplification of boson wave
- Create large bosonic cloud around spinning BHs
- Cloud decays through emission of long-duration quasi-monochromatic GWs ($f_{\text{GW}} \simeq mc^2/\pi\hbar$)

Black hole superradiance - vector field


M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019

...

New massive vector boson

$$\mathcal{L} \supset -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - \frac{1}{2}m^2 A'^{\mu}A'_{\mu}$$

$$\frac{h}{mc} \sim \frac{G_N M}{c^2} \quad \sim$$



- BH loses energy and angular momentum in favor of amplification of boson wave
- Create large bosonic cloud around spinning BHs
- Cloud decays through emission of long-duration quasi-monochromatic GWs ($f_{\text{GW}} \simeq mc^2/\pi\hbar$)


Allows to probe the existence of new ultralight fields, irrespectively of their cosmological production

Black hole superradiance - vector field

M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019


...

$$|\vec{E}'| \sim \alpha^{5/2} m M_{\text{pl}}$$



Credit: Nils Siemonsen

$$|\vec{B}'| \sim \alpha |\vec{E}'| \ll |\vec{E}'|$$




Black hole superradiance - vector field

M. Baryakhtar, R. Lasenby, M. Teo, PRD 2017
D. Bauman, H.S. Chia, J. Stout, L.T. Haar JCAP 2019


...

$$|\vec{E}'| \sim \alpha^{5/2} m M_{\text{pl}}$$



Credit: Nils Siemonsen


$$|\vec{B}'| \sim \alpha |\vec{E}'| \ll |\vec{E}'|$$



Kinetically-mixed dark photon cloud

N. Siemonsen, CM, D. Egaña-Ugrinovic, J. Huang, M. Baryakhtar, W. E. East
Phys. Rev. D (2023), arxiv:2212.09772


Cloud of large rotating electric field



Kinetically-mixed dark photon cloud


N. Siemonsen, CM, D. Egaña-Ugrinovic, J. Huang, M. Baryakhtar, W. E. East
Phys. Rev. D (2023), arxiv:2212.09772

Production of charged particles:
screening




Rotating dipole

N. Siemonsen, CM, D. Egaña-Ugrinovic, J. Huang, M. Baryakhtar, W. E. East
Phys. Rev. D (2023), arxiv:2212.09772




Kinetically-mixed dark photon cloud

N. Siemonsen, CM, D. Egaña-Ugrinovic, J. Huang, M. Baryakhtar, W. E. East
Phys. Rev. D (2023), arxiv:2212.09772



For the case with accretion disk, see:
S. Xin and E. R. Most, Phys.Rev.D 111 (2025), arxiv:2406.02992


Superradiance cloud + plasma




?

Neutron star pulsar

Magnetosphere of neutron star pulsar





Emission spectra of the Vela pulsar



From *Pulsar Magnetospheres and Their Radiation*

A. Philippov and M. Kramer, *Annu. Rev. Astron. Astrophys.* 2022


Dark photon cloud-pulsar analogies




- Large EM fields
- Rotating objects ($\sim$ ms periods)

Dark photon cloud-pulsar differences

$$|\vec{B}'| \sim \alpha |\vec{E}'| \ll |\vec{E}'|$$



Magnetically dominated




Electrically dominated



Dissipation is important

Superradiance cloud + plasma modeling



Nils Siemonsen



Full GR numerical MHD simulations


$$\partial_t \vec{B} = -\nabla \times \vec{E}$$

$$\partial_t \vec{E} = \nabla \times \vec{B} - \vec{J} - \epsilon m^2 \vec{A}'$$

Ohm's law $\vec{J} = \sigma(\vec{E} + \vec{v} \times \vec{B})$

- Solve equations for the E, B fields
- Compute EM energy emitted for $\sigma \rightarrow \infty$ ($\sigma \gg m$)


Simulations: B field & energy dissipation



Nils Siemonsen




The dissipative energy density is maximal in the regions of magnetic field lines reconnection.




Magnetic reconnection

Plasma density




- Discontinuity in the magnetic field lines
- In a **neutron star pulsars** it happens on the equatorial plane (2d structures “current sheets”)



Field energy dissipated into particle acceleration and high energy emissions

Dark photon cloud's luminosity

Numerical solutions and fit




- Poynting flux

$$P_{EM} = \oint dS \vec{r} \cdot (\vec{E} \times \vec{B})$$


- Dissipative losses

$$L_{diss} = \int dV \vec{E} \cdot \vec{J}$$



Dark photon cloud's luminosity

N. Siemonsen, CM, D. Egaña-Ugrinovic, J. Huang, M. Baryakhtar, W. E. East
Phys. Rev. D (2023), arxiv:2212.09772



$\epsilon \ll 1$

γ γ'



$$L_{\text{fit}} = \epsilon^2 F(\alpha) \frac{M_c}{GM} \simeq 10^{43} \text{ erg/s}$$

Crab pulsar $L_{\text{crab}} \simeq 10^{38} \text{ erg/s}$

Supernova $L_{\text{SN}} \simeq 10^{43} - 10^{45} \text{ erg/s}$


“New pulsar”: periodicity and spectrum

- Cannot be robustly inferred from current simulations
- Pulsar analogy: low energy periodic component + X-ray/ γ -ray emissions



Multimessenger searches


“New pulsar” signals




- GW emissions with frequency $mc^2/\pi\hbar \sim \text{few} \times 100 \text{ Hz}$
- Low and high energy EM emissions with periodicity $2\pi\hbar/mc^2 \sim \text{ms}$
- Decays over time with timescales from $\sim \text{days}$ to $\sim \text{thousands of years}$

Multimessenger signatures

1. EM follow-ups of compact binary mergers
(large $\alpha \equiv G_N M m$, young system)




2. GW follow-ups of anomalous pulsars
(small $\alpha \equiv G_N M m$, old system)




Multimessenger signatures

1. EM follow-ups of compact binary mergers
(large $\alpha \equiv G_N M m$, young system)



2. GW follow-ups of anomalous pulsars
(small $\alpha \equiv G_N M m$, old system)



Source selection


3 classes of selected sources:

- Spinning-up $\dot{f}_{\text{rot}} > 0$
- Frequency doublets: 2 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$
- Frequency triplets: 3 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$

Source selection

3 classes of selected sources:

- Spinning-up $\dot{f}_{\text{rot}} > 0$
- Frequency doublets: 2 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$
- Frequency triplets: 3 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$




59 sources selected
from ATNF pulsar catalog

Source selection

3 classes of selected sources:


- Spinning-up $\dot{f}_{\text{rot}} > 0$
- Frequency doublets: 2 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$
- Frequency triplets: 3 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$



Source selection

3 classes of selected sources:

- Spinning-up $\dot{f}_{\text{rot}} > 0$
- Frequency doublets: 2 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$
- Frequency triplets: 3 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq 10^{-3}$



Source selection


3 classes of selected sources:

- Spinning-up $\dot{f}_{\text{rot}} > 0$

- Frequency doublets: 2 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq$

- Frequency triplets: 3 pulsars with $\frac{|f_i - f_j|}{f_i} < \frac{v_{\text{rad}}}{c} \simeq$

- ▲ Narrow-band
- ◆ Targeted
- Semi-coherent
- + All-sky
- Binary
- × Galactic Center
- ⊙ Sun



15 discarded

(binary systems w/ bad ephemerides) **Used to look for dark photon signals**

➡ 59 sources selected
from ATNF pulsar catalog

44 sources

4 sources already targeted LVK O2-O3

R. Abbott et al., Astrophys.J. 935, 1 (2022), arxiv:2111.13106

6 w/ bad sky position measurements ->
using LVK O3 all-sky search

R. Abbott et al., Phys.Rev.D. 106, 102008 (2022), arxiv:2201.00697

34 analyzed for the first time

Methods for the analysis

Lorenzo Mirasola, Francesco Amicucci,
Sabrina D'Antonio, Cristiano Palomba,
Paola Leaci, Luca D'Onofrio, Pia Astone

Isolated sources

Maximum coherence time insensitive to
sky position uncertainty:

$$T_{\text{FFT}}^{\text{sky}}(\delta\beta) \approx \frac{10^4}{f_0 \delta\beta \sin \beta}$$

$$T_{\text{FFT}}^{\text{sky}}(\delta\lambda) \approx \frac{10^4}{f_0 \delta\lambda \cos \beta}$$

From Mirasola+ (2024)

Methods for the analysis

Lorenzo Mirasola, Francesco Amicucci,
Sabrina D'Antonio, Cristiano Palomba,
Paola Leaci, Luca D'Onofrio, Pia Astone


Isolated sources

Maximum coherence time insensitive to
sky position uncertainty:

$$T_{\text{FFT}}^{\text{sky}}(\delta\beta) \approx \frac{10^4}{f_0 \delta\beta \sin \beta}$$

$$T_{\text{FFT}}^{\text{sky}}(\delta\lambda) \approx \frac{10^4}{f_0 \delta\lambda \cos \beta}$$

From Mirasola+ (2024)



$$T_{\text{FFT}}^{\text{sky}} \gg T_{\text{obs}}$$

5-vector narrowband
Mastrogiovanni+ CQG (2017)

$$T_{\text{FH}} < T_{\text{FFT}}^{\text{sky}} \lesssim T_{\text{obs}}$$

5-vector semi-coherent
D'Antonio+ PRD (2023)

$$T_{\text{FH}} \sim T_{\text{FFT}}^{\text{sky}} \ll T_{\text{obs}}$$

results from all-sky Frequency-Hough
Abbott+, PRD (2022)

Methods for the analysis

Lorenzo Mirasola, Francesco Amicucci,
Sabrina D'Antonio, Cristiano Palomba,
Paola Leaci, Luca D'Onofrio, Pia Astone


Isolated sources

Maximum coherence time insensitive to
sky position uncertainty:

$$T_{\text{FFT}}^{\text{sky}}(\delta\beta) \approx \frac{10^4}{f_0 \delta\beta \sin \beta}$$

$$T_{\text{FFT}}^{\text{sky}}(\delta\lambda) \approx \frac{10^4}{f_0 \delta\lambda \cos \beta}$$

From Mirasola+ (2024)



$$T_{\text{FFT}}^{\text{sky}} \gg T_{\text{obs}}$$

5-vector narrowband
Mastrogiovanni+ CQG (2017)

$$T_{\text{FH}} < T_{\text{FFT}}^{\text{sky}} \lesssim T_{\text{obs}}$$

5-vector semi-coherent
D'Antonio+ PRD (2023)

$$T_{\text{FH}} \sim T_{\text{FFT}}^{\text{sky}} \ll T_{\text{obs}}$$

results from all-sky Frequency-Hough
Abbott+, PRD (2022)

Binary sources

5-vector for sources in a binary system
Singhal+ CQG (2019)

Semi-coherent for sources in a binary system
Mirasola+ (2024)

Methods for the analysis

Lorenzo Mirasola, Francesco Amicucci,
Sabrina D'Antonio, Cristiano Palomba,
Paola Leaci, Luca D'Onofrio, Pia Astone

Isolated sources

Maximum coherence time insensitive to
sky position uncertainty:

$$T_{\text{FFT}}^{\text{sky}}(\delta\beta) \approx \frac{10^4}{f_0 \delta\beta \sin \beta}$$

$$T_{\text{FFT}}^{\text{sky}}(\delta\lambda) \approx \frac{10^4}{f_0 \delta\lambda \cos \beta}$$

From Mirasola+ (2024)

$$T_{\text{FFT}}^{\text{sky}} \gg T_{\text{obs}}$$

5-vector narrowband
Mastrogiovanni+ CQG (2017)

$$T_{\text{FH}} < T_{\text{FFT}}^{\text{sky}} \lesssim T_{\text{obs}}$$

5-vector semi-coherent
D'Antonio+ PRD (2023)

$$T_{\text{FH}} \sim T_{\text{FFT}}^{\text{sky}} \ll T_{\text{obs}}$$

results from all-sky Frequency-Hough
Abbott+, PRD (2022)

Binary sources

5-vector for sources in a binary system
Singhal+ CQG (2019)

Semi-coherent for sources in a binary system
Mirasola+ (2024)

False alarm probability for outliers at 1%



Credits: Lorenzo Mirasola

Results analysis with O3 data

L. Mirasola, **CM**, et al.
Phys. Rev. D 11 (2025), arxiv:2501.02052


No outliers identified



Upper limits at the
95% confidence level




ULs below the maximum
predicted strain



Interpretation of CW strain ULs

CW strain at the detector:

$$h_0(M, \chi, \tau_{\text{obs}}, d|m) = \frac{h_0^s(M, \chi|m)}{1 + \tau_{\text{obs}}/\tau_{\text{GW}}(M, \chi|m)} \frac{1}{d},$$




Interpretation of CW strain ULs

CW strain at the detector:

$$h_0(M, \chi, \tau_{\text{obs}}, d|m) = \frac{h_0^s(M, \chi|m)}{1 + \tau_{\text{obs}}/\tau_{\text{GW}}(M, \chi|m)} \frac{1}{d},$$

Dark photon
 m




Fixed by the radio observation

$$f_{\text{GW}} = 2f_{\text{rot}} = mc^2/\pi\hbar$$

BH properties


$M, \chi, \tau, d,$
of BHs




Integrate over,
Given a BH population

BH in the Galactic disk


BH mass distribution




BH spin distribution



BH age distribution



BH spatial distribution
(galactic disk)




Interpretation of CW strain ULs

CW strain at the detector:

$$h_0(M, \chi, \tau_{\text{obs}}, d|m) = \frac{h_0^s(M, \chi|m)}{1 + \tau_{\text{obs}}/\tau_{\text{GW}}(M, \chi|m)} \frac{1}{d},$$

Dark photon
 m, ϵ




Fixed by the radio observation

$$f_{\text{GW}} = 2f_{\text{rot}} = mc^2/\pi\hbar$$

BH properties


$M, \chi, \tau, d,$
of BHs




Targeted search also depends on
EM dark photon coupling!

- Large enough for the cloud to be luminous
- Small enough so that the cloud does not decay too fast


BH mass distribution




BH spin distribution



BH age distribution



BH spatial distribution
(galactic disk)



Expected signal

$$\frac{dn_h}{d \log h_0}(h_0|m, \varepsilon) = \frac{x_{\text{disk}} N_{\text{BH}}}{2\pi \mathcal{N}} \int dM \frac{M_{\text{min}}^{1.35}}{M^{2.35}} \int d\chi \frac{1}{\chi_{\text{max}}} \frac{h_{r_d}}{h_0} \frac{\tau_{\text{GW}}}{\tau_{\text{max}}} \Theta_{\text{pl}} \int d\varphi \int d\tilde{\rho} \frac{\tilde{\rho} e^{-\tilde{\rho}}}{\sqrt{\tilde{\rho}^2 + \tilde{r}_{\text{obs}}^2 - 2\tilde{\rho} \tilde{r}_{\text{obs}} \cos \varphi}} \Theta_f \Theta_{\text{EM}} \Theta_{\text{ev}},$$

Expected signal

$$\frac{dn_h}{d \log h_0}(h_0|m, \varepsilon) = \frac{x_{\text{disk}} N_{\text{BH}}}{2\pi \mathcal{N}} \int dM \frac{M_{\text{min}}^{1.35}}{M^{2.35}} \int d\chi \frac{1}{\chi_{\text{max}}} \frac{h_{r_d}}{h_0} \frac{\tau_{\text{GW}}}{\tau_{\text{max}}} \Theta_{\text{pl}} \int d\varphi \int d\tilde{\rho} \frac{\tilde{\rho} e^{-\tilde{\rho}}}{\sqrt{\tilde{\rho}^2 + \tilde{r}_{\text{obs}}^2 - 2\tilde{\rho} \tilde{r}_{\text{obs}} \cos \varphi}} \Theta_f \Theta_{\text{EM}} \Theta_{\text{ev}},$$


$$N_{\text{ev}}(h_0 > h_0^{95\%} \mid m, \varepsilon) = \int_{h_0^{95\%}}^{\infty} dh_0 \frac{1}{h_0} \frac{dn_h}{d \log h_0}$$

Expected signal

$$\frac{dn_h}{d \log h_0}(h_0|m, \varepsilon) = \frac{x_{\text{disk}} N_{\text{BH}}}{2\pi \mathcal{N}} \int dM \frac{M_{\text{min}}^{1.35}}{M^{2.35}} \int d\chi \frac{1}{\chi_{\text{max}}} \frac{h_{r_d}}{h_0} \frac{\tau_{\text{GW}}}{\tau_{\text{max}}} \Theta_{\text{pl}} \int d\varphi \int d\tilde{\rho} \frac{\tilde{\rho} e^{-\tilde{\rho}}}{\sqrt{\tilde{\rho}^2 + \tilde{r}_{\text{obs}}^2 - 2\tilde{\rho} \tilde{r}_{\text{obs}} \cos \varphi}} \Theta_f \Theta_{\text{EM}} \Theta_{\text{ev}},$$

$$N_{\text{ev}}(h_0 > h_0^{95\%} | m, \varepsilon) = \int_{h_0^{95\%}}^{\infty} dh_0 \frac{1}{h_0} \frac{dn_h}{d \log h_0}$$


Signal strain distribution





$\varepsilon = 10^{-8}$

Fraction of EM power into radio:

Number of expected events




Number of expected events: dark photon parameter space



Fraction of EM power into radio: 10^{-5}

Several events expected above detection threshold

Number of expected events: dark photon parameter space



Several events expected above detection threshold

Depend on (not so well known):


- BH population properties
- Dark photon cloud EM emission properties

Not a robust “exclusion bound”, but a signal could have been seen.

Fraction of EM power into radio: 10^{-5}

Summary

- BH superradiance allows to probe the existence of new ultralight vector fields, irrespectively of their cosmological production
- Dark photons with EM coupling can lead to “new pulsars”
- Can be targeted with multimessenger searches



CW emission

EM emission

CW+EM

- CW searches targeting EM sources

BH merger event

EM emission


GW+EM

- EM follow-up of merger events

Summary

- BH superradiance allows to probe the existence of new ultralight vector fields, irrespectively of their cosmological production
- Dark photons with EM coupling can lead to “new pulsars”
- Can be targeted with multimessenger searches

Thank you!



CW emission

EM emission

CW+EM

- CW searches targeting EM sources

BH merger event



EM emission

GW+EM


- EM follow-up of merger events

Back-up

Dark photon superradiance parameter space



Simulation: periodicity?



- Fundamental time scale of the system $2\pi/\mu$
- Weak evidence of periodicity from our simulation
- Pulsar analogy

Simulation: spectrum?



- Synchrotron emission of highly boosted particles

$e^{\pm}$ boosted to $\gamma \sim \mathcal{O}(10^3)$

synchrotron photons $\sim$ few keV to MeV

- Coherent low-energy radio emission (continuous + periodic) ?



Expected signals



$$\varepsilon = 10^{-8}$$

Fraction of EM power into radio: 10^{-5}



Expected signals



$$\varepsilon = 10^{-8}$$

Fraction of EM power into radio: 10^{-3}


Expected signals



$$\varepsilon = 10^{-8}$$


Fraction of EM power into radio: 10^{-7}

Expected radio signals





$$\varepsilon = 10^{-8}$$

Fraction of EM power into radio: 10^{-5}



Detection threshold for large-scale radio pulsar surveys

Number of expected events



Fraction of EM power into radio: 10^{-5}

Observational signatures

BH superradiance can be used to probe the existence of ultralight bosons

0905.4720, String Axiverse

A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, J. March–Russell

1004.3558, Exploring the String Axiverse with Precision Black Hole Physics

A. Arvanitaki, S. Dubovsky

1411.2263, Discovering the QCD Axion with Black Holes and Gravitational Waves

A. Arvanitaki, M. Baryakhtar, X. Huang

1704.04791, Superradiant Instability and Backreaction of Massive Vector Fields around Kerr Black Holes

W. E. East, F. Pretorius

1801.01420, Constraining the mass of dark photons and axion–like particles through black–hole superradiance

V. Cardoso, O. J. C. Dias, G. S. Hartnett, M. Middleton, P. Pani, J. E. Santos

....

1. Spin distribution of BH population



2. Continuous gravitational wave signal



Observational signatures: spin distribution

BH superradiance can be used to probe the existence of ultralight bosons



Axion masses around 10^{-13} eV already disfavored!



* depends on the time between BH formation and the merger (longer merger time, longer time for SR to grow)

Observational signatures: continuous GW

BH superradiance can be used to probe the existence of ultralight bosons



Vector boson clouds could be seen with targeted CW searches of BH merger remnants in future LVK observing runs!

* needs good sky localization