

PRIMORDIAL BLACK HOLES, CHARGE
AND DARK MATTER:

RETHINKING EVAPORATION LIMITS


IN COLLABORATION WITH JUSTIN FENG, SEBASTIAN SCHUSTER,
AND MATT VISSER

PHYS. REV. D 112, 123529

JESSICA SANTIAGO

CENTRE DE PHYSIQUE THEORIQUE
AIX MARSEILLE UNIVERSITY

Primordial black holes as DM candidate



Carr, B., Kohri, K., Sendouda, Y. & Yokoyama, J. (2021)




Reissner Nordstrom metric

$$ds^2 = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2$$

- It is static and spherically symmetric;
- Solves the coupled Einstein-Maxwell equations
- Has two event-horizons located at:

$$r_{\pm} = M \pm \sqrt{M^2 - Q^2}$$



The temperature of an evaporating RN black hole is given by:

$$T = \frac{\hbar\kappa}{2\pi}, \quad \text{with} \quad \kappa = \frac{(M^2 - Q^2)^{1/2}}{r_+^2}$$

For standard EM:

$$\frac{Q}{M} \ll \frac{m}{e} \simeq 10^{-21} \quad (\text{in geometric units})$$



Reissner Nordstrom PBH with dark electric charge

$$ds^2 = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2$$

$$T = \frac{\hbar\kappa}{2\pi}, \quad \text{with} \quad \kappa = \frac{(M^2 - Q^2)^{1/2}}{r_+^2}$$



Dark electromagnetic charge

$$\mathcal{L} = \bar{\chi}(i\not{D} + m_\chi)\chi - \frac{1}{4}\hat{F}_{\mu\nu}\hat{F}^{\mu\nu}$$

$D_\mu = \partial_\mu - ie_\chi \hat{A}_\mu$ is the gauge covariant derivative

$\hat{F}_{\mu\nu}$ is the field-strength tensor for dark electromagnetism


Ackerman, L., Buckley, M. R., Carroll, S. M. & Kamionkowski, M. (2009)

Charged Black Hole Evaporation:


Two effects are responsible:

Hawking effect

Quantum field theory in curved spacetime effect. It is generated by the presence of an event horizon, which changes the notion of an initial vacuum state into a thermal bath.




$$\frac{dM}{dt} = -aT^4\alpha\sigma_0$$



Schwinger effect

The presence of a sufficiently strong electric field stimulates the quantum creation of pairs of particles.



The Schwinger expression for the rate of electron-positron pair creation per unit four-volume is given by:

$$\Gamma = \frac{(eE)^2}{4\pi^3\hbar^2c} \sum_{n=1}^{\infty} \frac{1}{n^2} \exp\left(-\frac{\pi m_e^2 c^3 n}{\hbar e E}\right)$$

HW assumptions -- Dark EM extension

Positive charge

Applicability of
Schwinger's result:

$$\Gamma = \frac{(eE)^2}{4\pi^3 \hbar^2 c} \sum_{n=1}^{\infty} \frac{1}{n^2} \exp\left(-\frac{\pi m_e^2 c^3 n}{\hbar e E}\right)$$

Black hole mass must be much larger than the reduced Compton wavelength of the electron (or lightest charged particle)

$$M \gg \frac{\hbar}{m_e} \simeq 10^{-15} M_{\odot}$$

Series truncation:

$$\frac{e^3 Q}{m_e^2 r^2} \ll 1$$

Error function series
truncation:

$$r_+^2 \gg QQ_0$$

Summarized model limitations:

$$(i) \ M \gg \frac{\hbar}{m_e}, \quad (ii) \ \frac{e^3 Q}{m_e^2 r^2} \ll 1, \quad \text{and} \quad (iii) \ r_+^2 \gg QQ_0$$

Dark EM extension:

$$M_{DE} \gg \frac{\hbar}{m_{\chi}} = \frac{10^{-15}}{\sigma_m} M_{\odot}$$

$$M_{DE} \gg \frac{e_{\chi}^3}{m_{\chi}^2} = 4 \cdot 10^3 \frac{\sigma_e^3}{\sigma_m^2} M_{\odot}$$

$$\frac{\sigma_e^2 \sigma_M}{\sigma_m} \gg 10^{-66}$$

RN BH Evaporation:

The full evolution equations:

$$\frac{dM}{dt} = -aT^4\alpha\sigma_0 + \frac{Q}{r_+} \frac{dQ}{dt},$$


$$\frac{dQ}{dt} = -\frac{e^3}{\pi^2\hbar^2 r_+} \exp\left(-\frac{r_+^2}{QQ_0}\right) - \frac{\pi}{\sqrt{QQ_0}} \operatorname{erfc}\left(\frac{r_+}{\sqrt{QQ_0}}\right)$$

$$\sigma_0 := \pi \frac{\left(3M + \sqrt{9M^2 - 8Q^2}\right)^4}{8\left(3M^2 - 2Q^2 + M\sqrt{9M^2 - 8Q^2}\right)}$$

$$Q_0 := \frac{\hbar e}{\pi m^2} \approx 1,7 \cdot 10^5 M_\odot$$

$$a = \pi^2 / (15\hbar^3)$$

Hiscock & Weems (1989)



RN BH Evaporation:

$$\frac{dM}{dt} = -aT^4\alpha\sigma_0 + \frac{Q}{r_+} \frac{dQ}{dt},$$

$$\frac{dQ}{dt} = -\frac{e^3}{\pi^2\hbar^2 r_+} \exp\left(-\frac{r_+^2}{QQ_0}\right) - \frac{\pi}{\sqrt{QQ_0}} \operatorname{erfc}\left(\frac{r_+}{\sqrt{QQ_0}}\right)$$

change of variables

$$Y := (Q/M)^2$$

$$\mu := M/M_s$$

$$\tau = \frac{s_0}{M_s} t$$

$$\frac{d\mu}{d\tau} = -\frac{(H(\mu, Y) + S(\mu, Y)Y^2)}{(\sqrt{1-Y} + 1)^4},$$


$$\frac{dY}{d\tau} = \frac{2(H(\mu, Y) - S(\mu, Y)(1 - Y + \sqrt{1-Y})Y)Y}{\mu(\sqrt{1-Y} + 1)^4}$$

$$H(\mu, Y) := \frac{(\sqrt{9-8Y} + 3)^4 (1-Y)^2}{\mu^2(\sqrt{1-Y} + 1)^4(3-2Y + \sqrt{9-8Y})},$$

$$S(\mu, Y) := \exp\left\{b_0\left[z_0 - \mu(\sqrt{1-Y} + 1)^2/\sqrt{Y}\right]\right\}.$$


the Hawking term

the Schwinger term




$$z_0 = \frac{e\hbar}{\pi m_e^2 M_s} \ln\left(\frac{960e^4 M_s^2}{\pi^2 \alpha m_e^2 \hbar^2}\right), \quad b_0 = \frac{m_e^2 M_s \pi}{e\hbar}$$


RN BH Evolution Profiles:




$$M = 0.3 \times 10^6 M_\odot, Y = 0.1$$




$$M = 9.0 \times 10^7 M_\odot, Y = 0.1$$



$$M = 1.50 \times 10^8 M_\odot, Y = 0.1$$



$$M = 1.68 \times 10^8 M_\odot, Y = 0.1$$




Approximate solutions

Charge dissipation zone


$$\frac{dY}{d\mu} = \frac{2S(\mu, Y) \left(\sqrt{1-Y} + 1 \right) Y^2}{\mu (H(\mu, Y) + S(\mu, Y) Y^2)} - \frac{2Y}{\mu}$$

Mass dissipation zone

$$Y_S(\mu) = \frac{(2\mu - \mu_1)\mu_1}{\mu^2}$$



$$Y_H(\mu) = \left(\frac{\mu_h}{\mu} \right)^2$$




Approximate solutions

$$\frac{dY}{d\mu} = \frac{2S(\mu, Y) \left(\sqrt{1-Y} + 1 \right) Y^2}{\mu (H(\mu, Y) + S(\mu, Y) Y^2)} - \frac{2Y}{\mu}$$

Approximate attractor curve

$$S(\mu, Y) := \exp \left\{ b_0 \left[z_0 - \mu \left(\sqrt{1-Y} + 1 \right)^2 / \sqrt{Y} \right] \right\}$$

$$z_0 - \frac{\mu \left(\sqrt{1-Y} + 1 \right)^2}{\sqrt{Y}} \quad \begin{cases} > 0 & \Rightarrow \text{Charge dissipation zone} \\ < 0 & \Rightarrow \text{Mass dissipation zone} \end{cases}$$



$$\mu = z_0 \frac{\sqrt{Y}}{(\sqrt{1-Y} + 1)^2}$$

Connection to Dark Matter

Rescaling relations

$$z_0^\chi = \frac{e_\chi \hbar}{\pi m_\chi^2 M_s} \ln \left(\frac{960 e_\chi^4 M_s^2}{\pi^2 \alpha m_\chi^2 \hbar^2} \right) \quad M_{z_0} = z_0 M_s$$

$$s_0^\chi = \frac{s_0}{\sigma_M^2}, \quad z_0^\chi = \frac{\sigma_e}{\sigma_m^2 \sigma_M} \left[z_0 + \frac{2}{b_0} \ln \left(\frac{\sigma_e^2 \sigma_M}{\sigma_m} \right) \right], \quad b_0^\chi = \frac{\sigma_m^2 \sigma_M}{\sigma_e} b_0$$

Varying the parameters

$$\begin{aligned} M_s^\chi &= \sigma_M M_s \\ m_\chi &= \sigma_m m_e \\ e_\chi &= \sigma_e e \end{aligned}$$

$$\xi = \sigma_e / \sigma_m$$

$$\sigma_e = \frac{\vartheta}{\sigma_M \xi}, \quad \sigma_m = \frac{\vartheta}{\sigma_M \xi^2}, \quad \xi = \xi(\vartheta) = \left(\frac{b_0 z_0 \vartheta}{b_0 z_0 + 2 \ln \vartheta} \right)^{1/3}$$



$$z_0^\chi = z_0 \quad M_{z_0}^\chi = \frac{\vartheta^{1/3}}{\sigma_m} \left(\frac{b_0 z_0 + 2 \ln \vartheta}{b_0 z_0} \right)^{2/3} M_{z_0}$$

Connection to Dark Matter


Varying the charge

$$z_0^\chi = \frac{e_\chi \hbar}{\pi m_\chi^2 M_s} \ln \left(\frac{960 e_\chi^4 M_s^2}{\pi^2 \alpha m_\chi^2 \hbar^2} \right).$$

Varying the mass



Decomposing the possible evolution scenarios



Small Y and/or small μ limit $(\mu \ll z_0)$


$$\frac{d\mu}{d\tau} = -\frac{(H(\mu, Y) + S(\mu, Y)Y^2)}{(\sqrt{1-Y} + 1)^4},$$

$$\Delta\tau \stackrel{Y \rightarrow 0}{=} -\int_{\mu}^0 \frac{32}{27} \tilde{\mu}^2 d\tilde{\mu}$$

$$\rightarrow \Delta\tau_{\text{schw}} \stackrel{Y \rightarrow 0}{=} \frac{32}{81} \mu^3$$

$$\Delta t = \frac{M_s^\chi}{s_0^\chi} \Delta\tau_{\text{schw}} = \frac{1920\pi}{\alpha\hbar} \left[\frac{32}{81} (M_s^\chi \mu)^3 \right]$$

$$= \frac{1920\pi}{\alpha\hbar} \left[\frac{32}{81} M^3 \right] = \Delta t_{\text{schw}}$$



Approximate attractor curve $(\mu = z_0)$

$$\mu = z_0 \frac{\sqrt{Y}}{(\sqrt{1-Y} + 1)^2}$$



$$S(\mu, Y) \sim 1$$






$$d\tau = -\frac{(\sqrt{1-Y} + 1)^4 d\mu}{(H(\mu, Y) + Y^2)}$$

$$\Delta\tau_{\text{att}}(\mu) = \frac{1}{a + b \mu^{-m}}$$

Approximate time estimate

$$M_{z_0} = M_s z_0$$

- $\mu < 0.05 \rightarrow (a, b, m) = (0,81/32,3)$
- $0.05 < \mu < 0.38 \rightarrow (a, b, m) = (-1.54,0.1045,4.765)$
- $\mu > 0.38 \rightarrow (a, b, m) = (2.308,0.0277,5.54)$








Near extremal evolution $(\mu > z_0)$




Near-extremal evolution:

$$\frac{d\mu}{d\tau} = -\exp[(z_0 - \mu)b_0]$$

$$\Delta t \sim \frac{M_s}{s_0^\chi b_0^\chi} \exp[(\mu_h - z_0^\chi) b_0^\chi]$$





New PBH mass bounds:



$$\Delta t_{\text{evap}}(\bar{\mu}) = \frac{M_s^\chi}{s_0^\chi b_0^\chi} \cdot \begin{cases} \Delta \tau_{\text{schw}}(\bar{\mu}) , \\ \Delta \tau_{\text{att}}(\bar{\mu}) , \\ (\exp[(\bar{\mu} - z_0^\chi) b_0^\chi] - 1) + \Delta \tau_{\text{att}}(z_0) \\ \exp[(\bar{\mu} - z_0^\chi) b_0^\chi] , \end{cases}$$

$$M_{\text{univ}} = \begin{cases} M_{\text{schw}} , & \bar{\mu} \ll z_0^\chi \\ M_{\text{att}}^{\text{univ}} , & \bar{\mu} \leq z_0^\chi \\ M_{\text{near-extr}}^{\text{univ}} , & \bar{\mu} > z_0^\chi \\ M_{\text{extr}} , & \bar{\mu} - z_0^\chi \gg 1/b_0^\chi \end{cases}$$


Long Live (tiny) Black Holes!



Conclusions

- Reissner-Nordstrom black holes have very interesting evaporation evolution for certain mass ranges;
- If PBHs can form with an initial dark U(1) charge, they would not discharge via accretion effects, allowing them to follow the mass dissipation curve and achieve near-extremality;
- Given the right combination of dark-electron charge, mass and initial PBH mass, PBHs of masses below $10^{(-15)}$ solar masses can still account for the total amount of dark matter in the universe.

- And Long Live Black Holes!
- Phys. Rev. D 112, 123529



The Schwinger effect validity region

Electric field gradients must be small compared to the Compton wavelength of the lightest electrically charged particle


$$\frac{1}{E} \frac{dE}{d(\text{proper length})} < \frac{1}{\lambda_{\text{compDE}}} = \frac{m_\chi}{\hbar c}.$$

$$\sqrt{1 - \frac{2M}{r} + \frac{Q^2}{r^2}} \frac{2}{r} < \frac{1}{\lambda_{\text{compDE}}} = \frac{m_\chi}{m_{\text{Planck}}^2}$$

$$r^4 - 4\lambda_{\text{compDE}}^2(r^2 - 2Mr + Q^2) > 0$$

$$r > r_+$$

$$r^2 - 2Mr + Q^2 > 0$$



Regarding the discreteness of states near extremality

Breaking down energy

$$M - Q \lesssim E_{\text{brk.}} \equiv \frac{M_{Pl}}{Q^3}.$$

(from 2411.03447 in weird units)

$$M - Q \gg m_{\text{Planck}}^4 / Q^3$$


$$Y^{3/2}(1 - \sqrt{Y}) \gg (m_{\text{Planck}}/M)^4$$

In our analysis

smallest Y value $\sqrt{Y} \sim 1 - 10^{-25}$

smallest mass $10^{-25} M_{\odot} \sim 10^{13} m_{\text{Planck}}$

$$(m_{\text{Planck}}/M)^4 \sim 10^{-52}$$



Near extremal evolution

Evaporation times

ϱ/ϱ_0	$\mu_h = 0.70$	$\mu_h = 0.75$	$\mu_h = 0.80$	$\mu_h = 0.85$	$\mu_h = 0.90$
10^{-14}	5.58×10^{31}	2.90×10^{36}	1.51×10^{41}	7.83×10^{45}	4.07×10^{50}
10^{-15}	4.94×10^{23}	6.82×10^{27}	9.42×10^{31}	1.30×10^{36}	1.80×10^{40}
10^{-16}	4.16×10^{15}	1.51×10^{19}	5.54×10^{22}	2.03×10^{26}	7.43×10^{29}
10^{-17}	3.29×10^7	3.12×10^{10}	3.00×10^{13}	2.90×10^{16}	2.80×10^{19}
10^{-18}	2.43×10^{-1}	5.80×10^1	1.45×10^4	3.64×10^6	9.20×10^8

Table 1: Evaporation times (in years) as a function of the ‘hanging mass’ μ_h and charge to mass ratio, where $\varrho = e_\chi/m_\chi$ is the rescaled charge to mass ratio, and $\varrho_0 = 2.04 \times 10^{21} \sqrt{4\pi\epsilon_0 G}$ corresponds to the standard value in electrodynamics. In each case, the electron charge is rescaled by a factor $\sigma_e = 10^{-4}$.

Evaporation times: approximation vs. numerical results

ϱ/ϱ_0	$\mu_h = 0.70$	$\mu_h = 0.75$	$\mu_h = 0.80$	$\mu_h = 0.85$	$\mu_h = 0.90$
10^{-14}	4.83×10^{-6}	1.26×10^{-7}	3.34×10^{-9}	8.99×10^{-11}	3.25×10^{-12}
10^{-15}	2.52×10^{-5}	1.00×10^{-6}	4.05×10^{-8}	1.65×10^{-9}	6.90×10^{-11}
10^{-16}	1.46×10^{-4}	8.67×10^{-6}	5.28×10^{-7}	3.28×10^{-8}	2.06×10^{-9}
10^{-17}	1.13×10^{-3}	9.25×10^{-5}	8.13×10^{-6}	7.45×10^{-7}	7.03×10^{-8}
10^{-18}	4.98×10^{-2}	3.12×10^{-3}	2.33×10^{-4}	2.59×10^{-5}	3.31×10^{-6}

