

The gauge-invariant parametrized post-Newtonian formalism

arXiv:1910.09245

Manuel Hohmann

Laboratory of Theoretical Physics - Institute of Physics - University of Tartu
Center of Excellence "The Dark Side of the Universe"



Gravitational waves, black holes and fundamental physics - 15. 1.
2020

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations
- 3 Parametrized post-Newtonian formalism
- 4 Gauge-invariant PPN formalism
- 5 Example: PPN limit of scalar-tensor gravity
- 6 Conclusion

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations
- 3 Parametrized post-Newtonian formalism
- 4 Gauge-invariant PPN formalism
- 5 Example: PPN limit of scalar-tensor gravity
- 6 Conclusion

- Experimental tests of modified gravity theories:
 - Cosmological observations (CMB, supernovae, ...).
 - Gravitational waves.
 - Direct observation of black holes.
 - Solar system, pulsars, ...

Motivation

- Experimental tests of modified gravity theories:
 - Cosmological observations (CMB, supernovae, ...).
 - Gravitational waves.
 - Direct observation of black holes.
 - **Solar system, pulsars, ...**
- Parametrized post-Newtonian formalism:
 - Weak-field approximation of metric gravity theories.
 - Assumes particular coordinate system (“universe rest frame”).
 - Characterizes gravity theories by 10 (constant) parameters.
 - Parameters closely related to solar system observations.

- Experimental tests of modified gravity theories:
 - Cosmological observations (CMB, supernovae, ...).
 - Gravitational waves.
 - Direct observation of black holes.
 - Solar system, pulsars, ...
- Parametrized post-Newtonian formalism:
 - Weak-field approximation of metric gravity theories.
 - Assumes particular coordinate system (“universe rest frame”).
 - Characterizes gravity theories by 10 (constant) parameters.
 - Parameters closely related to solar system observations.
- ↪ Extensions of the PPN formalism:
 - More fundamental fields constituting the metric.
 - More than one dynamical metric.
 - Diffeomorphism invariant / purely geometric formalism.

- Experimental tests of modified gravity theories:
 - Cosmological observations (CMB, supernovae, ...).
 - Gravitational waves.
 - Direct observation of black holes.
 - Solar system, pulsars, ...
- Parametrized post-Newtonian formalism:
 - Weak-field approximation of metric gravity theories.
 - Assumes particular coordinate system (“universe rest frame”).
 - Characterizes gravity theories by 10 (constant) parameters.
 - Parameters closely related to solar system observations.
- ↪ Extensions of the PPN formalism:
 - More fundamental fields constituting the metric.
 - More than one dynamical metric.
 - Diffeomorphism invariant / purely geometric formalism.
- ↪ Use gauge-invariant higher order perturbation theory.

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations**
- 3 Parametrized post-Newtonian formalism
- 4 Gauge-invariant PPN formalism
- 5 Example: PPN limit of scalar-tensor gravity
- 6 Conclusion

Concept and use of gauge

- Reference spacetime:
 - Manifold M_0 with metric $g^{(0)}$ and coordinates (x^μ) .
 - Usually some highly symmetric standard spacetime:
 - maximally symmetric spacetime: Minkowski, (anti-)de Sitter
 - cosmological (background) solution of a gravity theory

Concept and use of gauge

- Reference spacetime:
 - Manifold M_0 with metric $g^{(0)}$ and coordinates (x^μ) .
 - Usually some highly symmetric standard spacetime:
 - maximally symmetric spacetime: Minkowski, (anti-)de Sitter
 - cosmological (background) solution of a gravity theory
- Physical spacetime:
 - Manifold M with metric g .
 - No preferred / canonical choice of coordinates.

Concept and use of gauge

- Reference spacetime:
 - Manifold M_0 with metric $g^{(0)}$ and coordinates (x^μ) .
 - Usually some highly symmetric standard spacetime:
 - maximally symmetric spacetime: Minkowski, (anti-)de Sitter
 - cosmological (background) solution of a gravity theory
 - Physical spacetime:
 - Manifold M with metric g .
 - No preferred / canonical choice of coordinates.
- ⚡ No canonical relation between physical and reference spacetime.

Concept and use of gauge

- Reference spacetime:
 - Manifold M_0 with metric $g^{(0)}$ and coordinates (x^μ) .
 - Usually some highly symmetric standard spacetime:
 - maximally symmetric spacetime: Minkowski, (anti-)de Sitter
 - cosmological (background) solution of a gravity theory
 - Physical spacetime:
 - Manifold M with metric g .
 - No preferred / canonical choice of coordinates.
- ⚡ No canonical relation between physical and reference spacetime.
1. No identification of points on M and M_0 : no coordinates on M .

Concept and use of gauge

- Reference spacetime:
 - Manifold M_0 with metric $g^{(0)}$ and coordinates (x^μ) .
 - Usually some highly symmetric standard spacetime:
 - maximally symmetric spacetime: Minkowski, (anti-)de Sitter
 - cosmological (background) solution of a gravity theory
- Physical spacetime:
 - Manifold M with metric g .
 - No preferred / canonical choice of coordinates.
- ⚡ No canonical relation between physical and reference spacetime.
 1. No identification of points on M and M_0 : no coordinates on M .
 2. No possibility to compare g and $g^{(0)}$: different manifolds.

Concept and use of gauge

- Reference spacetime:
 - Manifold M_0 with metric $g^{(0)}$ and coordinates (x^μ) .
 - Usually some highly symmetric standard spacetime:
 - maximally symmetric spacetime: Minkowski, (anti-)de Sitter
 - cosmological (background) solution of a gravity theory
- Physical spacetime:
 - Manifold M with metric g .
 - No preferred / canonical choice of coordinates.
- ⚡ No canonical relation between physical and reference spacetime.
 1. No identification of points on M and M_0 : no coordinates on M .
 2. No possibility to compare g and $g^{(0)}$: different manifolds.
- ↪ Introduce a *gauge*: diffeomorphism $\mathcal{X} : M_0 \rightarrow M$.
 1. Identification of (coordinated) points on M and M_0 .
 2. Comparison between reference metric $g^{(0)}$ and ${}^{\mathcal{X}}g = \mathcal{X}^*g$ on M_0 .

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 .
- ⇒ Family of physical spacetimes (M_ϵ, g_ϵ) .

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 .
 - ⇒ Family of physical spacetimes (M_ϵ, g_ϵ) .
- Aim: perturbative expansion of g_ϵ in ϵ around $g^{(0)}$.

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 .

⇒ Family of physical spacetimes (M_ϵ, g_ϵ) .
- Aim: perturbative expansion of g_ϵ in ϵ around $g^{(0)}$.
- ⚡ Metrics g_ϵ are defined on different manifolds for different ϵ .

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 .

⇒ Family of physical spacetimes (M_ϵ, g_ϵ) .
- Aim: perturbative expansion of g_ϵ in ϵ around $g^{(0)}$.
- ⚡ Metrics g_ϵ are defined on different manifolds for different ϵ .
- ⇒ Use gauge to relate different manifolds:
 - Family of diffeomorphisms $\mathcal{X}_\epsilon : M_0 \rightarrow M_\epsilon$.
 - Assume $\mathcal{X}_0 = \text{id}_{M_0}$ on the reference spacetime.

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 .

⇒ Family of physical spacetimes (M_ϵ, g_ϵ) .
- Aim: perturbative expansion of g_ϵ in ϵ around $g^{(0)}$.
- ⚡ Metrics g_ϵ are defined on different manifolds for different ϵ .
- ⇒ Use gauge to relate different manifolds:
 - Family of diffeomorphisms $\mathcal{X}_\epsilon : M_0 \rightarrow M_\epsilon$.
 - Assume $\mathcal{X}_0 = \text{id}_{M_0}$ on the reference spacetime.
- Perturbative expansion:
 - Pullback $\mathcal{X}^* g_\epsilon = \mathcal{X}_\epsilon^* g_\epsilon$ defined on M_0 .

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 .
 - ⇒ Family of physical spacetimes (M_ϵ, g_ϵ) .
- Aim: perturbative expansion of g_ϵ in ϵ around $g^{(0)}$.
- ⚡ Metrics g_ϵ are defined on different manifolds for different ϵ .
- ⇒ Use gauge to relate different manifolds:
 - Family of diffeomorphisms $\mathcal{X}_\epsilon : M_0 \rightarrow M_\epsilon$.
 - Assume $\mathcal{X}_0 = \text{id}_{M_0}$ on the reference spacetime.
- Perturbative expansion:
 - Pullback ${}^{\mathcal{X}}g_\epsilon = \mathcal{X}_\epsilon^* g_\epsilon$ defined on M_0 .
 - Introduce series expansion in ϵ :

$${}^{\mathcal{X}}g_\epsilon = \sum_{k=0}^{\infty} \frac{\epsilon^k}{k!} \left. \frac{\partial^k {}^{\mathcal{X}}g_\epsilon}{\partial \epsilon^k} \right|_{\epsilon=0} = \sum_{k=0}^{\infty} \frac{\epsilon^k}{k!} {}^{\mathcal{X}}g^{(k)}.$$

Gauge and perturbations

- Parameter dependent physical metric:
 - Assume physical metric $g \equiv g_\epsilon$ depends on parameter $\epsilon \in \mathbb{R}$.
 - Assume every g_ϵ is defined on its own M_ϵ .
 - Assume $g_0 = g^{(0)}$ is the reference metric on M_0 . \Rightarrow Family of physical spacetimes (M_ϵ, g_ϵ) .

- Aim: perturbative expansion of g_ϵ in ϵ around $g^{(0)}$.

⚡ Metrics g_ϵ are defined on different manifolds for different ϵ .

\Rightarrow Use gauge to relate different manifolds:

- Family of diffeomorphisms $\mathcal{X}_\epsilon : M_0 \rightarrow M_\epsilon$.
- Assume $\mathcal{X}_0 = \text{id}_{M_0}$ on the reference spacetime.
- Perturbative expansion:
 - Pullback ${}^{\mathcal{X}}g_\epsilon = \mathcal{X}_\epsilon^* g_\epsilon$ defined on M_0 .
 - Introduce series expansion in ϵ :

$${}^{\mathcal{X}}g_\epsilon = \sum_{k=0}^{\infty} \frac{\epsilon^k}{k!} \left. \frac{\partial^k {}^{\mathcal{X}}g_\epsilon}{\partial \epsilon^k} \right|_{\epsilon=0} = \sum_{k=0}^{\infty} \frac{\epsilon^k}{k!} {}^{\mathcal{X}}g^{(k)}.$$

- Series coefficients ${}^{\mathcal{X}}g^{(k)}$ depend on gauge choice \mathcal{X} .

Gauge invariant perturbations

- Choose a fixed “distinguished” gauge $\mathcal{S}_\epsilon : M_0 \rightarrow M_\epsilon$:
 - E.g., impose gauge conditions on the metric.
 - Examples: harmonic gauge used for GWs, standard PPN gauge.

Gauge invariant perturbations

- Choose a fixed “distinguished” gauge $\mathcal{S}_\epsilon : M_0 \rightarrow M_\epsilon$:
 - E.g., impose gauge conditions on the metric.
 - Examples: harmonic gauge used for GWs, standard PPN gauge.
- Define gauge-invariant metric $\mathbf{g}_\epsilon = \mathcal{S} g_\epsilon = \mathcal{S}_\epsilon^* g_\epsilon$.

Gauge invariant perturbations

- Choose a fixed “distinguished” gauge $\mathcal{S}_\epsilon : M_0 \rightarrow M_\epsilon$:
 - E.g., impose gauge conditions on the metric.
 - Examples: harmonic gauge used for GWs, standard PPN gauge.
- Define gauge-invariant metric $\mathbf{g}_\epsilon = \mathcal{S}_\epsilon^* \mathbf{g}_\epsilon$.
- Metric in arbitrary gauge \mathcal{X} :

$${}^{\mathcal{X}}\mathbf{g}_\epsilon = \sum_{l_1=0}^{\infty} \cdots \sum_{l_k=0}^{\infty} \cdots \frac{\epsilon^{l_1+\cdots+kl_k+\cdots}}{(1!)^{l_1}\cdots(k!)^{l_k}\cdots l_1!\cdots l_k!\cdots} \mathbf{\xi}_{\mathcal{X}(1)}^{l_1} \cdots \mathbf{\xi}_{\mathcal{X}(k)}^{l_k} \cdots \mathbf{g}_\epsilon.$$

Gauge invariant perturbations

- Choose a fixed “distinguished” gauge $\mathcal{S}_\epsilon : M_0 \rightarrow M_\epsilon$:
 - E.g., impose gauge conditions on the metric.
 - Examples: harmonic gauge used for GWs, standard PPN gauge.
- Define gauge-invariant metric $\mathbf{g}_\epsilon = \mathcal{S}_\epsilon^* g_\epsilon = \mathcal{S}_\epsilon^* g_\epsilon$.
- Metric in arbitrary gauge \mathcal{X} :

$$\mathcal{X} g_\epsilon = \sum_{l_1=0}^{\infty} \cdots \sum_{l_k=0}^{\infty} \cdots \frac{\epsilon^{l_1+\cdots+kl_k+\cdots}}{(1!)^{l_1} \cdots (k!)^{l_k} \cdots l_1! \cdots l_k! \cdots} \mathcal{X}_{(1)}^{l_1} \cdots \mathcal{X}_{(k)}^{l_k} \cdots \mathbf{g}_\epsilon.$$

- Metric components split into two parts:
 - Gauge-invariant part \mathbf{g}_ϵ : physical content.
 - Gauge defining vector fields $\mathcal{X}_{(k)}$: coordinate choice.

Gauge invariant perturbations

- Choose a fixed “distinguished” gauge $\mathcal{S}_\epsilon : M_0 \rightarrow M_\epsilon$:
 - E.g., impose gauge conditions on the metric.
 - Examples: harmonic gauge used for GWs, standard PPN gauge.
- Define gauge-invariant metric $\mathbf{g}_\epsilon = \mathcal{S}^* \mathbf{g}_\epsilon = \mathcal{S}_\epsilon^* \mathbf{g}_\epsilon$.
- Metric in arbitrary gauge \mathcal{X} :

$${}^{\mathcal{X}}\mathbf{g}_\epsilon = \sum_{l_1=0}^{\infty} \cdots \sum_{l_k=0}^{\infty} \cdots \frac{\epsilon^{l_1+\cdots+kl_k+\cdots}}{(1!)^{l_1}\cdots(k!)^{l_k}\cdots l_1!\cdots l_k!\cdots} \mathbf{X}_{(1)}^{l_1} \cdots \mathbf{X}_{(k)}^{l_k} \cdots \mathbf{g}_\epsilon.$$

- Metric components split into two parts:
 - Gauge-invariant part \mathbf{g}_ϵ : physical content.
 - Gauge defining vector fields $\mathbf{X}_{(k)}$: coordinate choice.
- Number # of independent components:

$$\#({}^{\mathcal{X}}\mathbf{g}_\epsilon) = \#(\mathbf{g}_\epsilon) + \#(\mathbf{X}_{(k)}).$$

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations
- 3 Parametrized post-Newtonian formalism**
- 4 Gauge-invariant PPN formalism
- 5 Example: PPN limit of scalar-tensor gravity
- 6 Conclusion

Post-Newtonian matter and velocity orders

- Energy-momentum tensor of a perfect fluid:

$$T^{\mu\nu} = (\rho + \rho\Pi + p)u^\mu u^\nu + pg^{\mu\nu} .$$

- Rest mass density ρ .
- Specific internal energy Π .
- Pressure p .
- Four-velocity u^μ .

Post-Newtonian matter and velocity orders

- Energy-momentum tensor of a perfect fluid:

$$T^{\mu\nu} = (\rho + \rho\Pi + p)u^\mu u^\nu + pg^{\mu\nu}.$$

- Rest mass density ρ .
- Specific internal energy Π .
- Pressure p .
- Four-velocity u^μ .
- Universe rest frame and slow-moving source matter:
 - Consider some gauge $\mathcal{X} : M_0 \rightarrow M$ (“universe rest frame”).
 - Pullback of metric and matter variables along \mathcal{X} .
 - Velocity of the source matter: ${}^{\mathcal{X}}v^i = {}^{\mathcal{X}}u^i / {}^{\mathcal{X}}u^0$.
 - Assume that source matter is slow-moving: $|{}^{\mathcal{X}}\vec{v}| \ll 1$.

Post-Newtonian matter and velocity orders

- Energy-momentum tensor of a perfect fluid:

$$T^{\mu\nu} = (\rho + \rho\Pi + p)u^\mu u^\nu + pg^{\mu\nu}.$$

- Rest mass density ρ .
- Specific internal energy Π .
- Pressure p .
- Four-velocity u^μ .
- Universe rest frame and slow-moving source matter:
 - Consider some gauge $\mathcal{X} : M_0 \rightarrow M$ (“universe rest frame”).
 - Pullback of metric and matter variables along \mathcal{X} .
 - Velocity of the source matter: ${}^{\mathcal{X}}v^i = {}^{\mathcal{X}}u^i / {}^{\mathcal{X}}u^0$.
 - Assume that source matter is slow-moving: $|{}^{\mathcal{X}}\vec{v}| \ll 1$.
- Use $\epsilon = |{}^{\mathcal{X}}\vec{v}|$ as perturbation parameter.

Post-Newtonian matter and velocity orders

- Energy-momentum tensor of a perfect fluid:

$$T^{\mu\nu} = (\rho + \rho\Pi + p)u^\mu u^\nu + pg^{\mu\nu}.$$

- Rest mass density $\rho \sim \mathcal{O}(2)$.
- Specific internal energy $\Pi \sim \mathcal{O}(2)$.
- Pressure $p \sim \mathcal{O}(4)$.
- Four-velocity u^μ .
- Universe rest frame and slow-moving source matter:
 - Consider some gauge $\mathcal{X} : M_0 \rightarrow M$ (“universe rest frame”).
 - Pullback of metric and matter variables along \mathcal{X} .
 - Velocity of the source matter: ${}^{\mathcal{X}}v^i = {}^{\mathcal{X}}u^i / {}^{\mathcal{X}}u^0$.
 - Assume that source matter is slow-moving: $|{}^{\mathcal{X}}\vec{v}| \ll 1$.
- Use $\epsilon = |{}^{\mathcal{X}}\vec{v}|$ as perturbation parameter.
- Assign velocity orders $\mathcal{O}(n) \sim \epsilon^n$ to all quantities.

Post-Newtonian matter and velocity orders

- Energy-momentum tensor of a perfect fluid:

$$T^{\mu\nu} = (\rho + \rho\Pi + p)u^\mu u^\nu + pg^{\mu\nu}.$$

- Rest mass density $\rho \sim \mathcal{O}(2)$.
- Specific internal energy $\Pi \sim \mathcal{O}(2)$.
- Pressure $p \sim \mathcal{O}(4)$.
- Four-velocity u^μ .
- Universe rest frame and slow-moving source matter:
 - Consider some gauge $\mathcal{X} : M_0 \rightarrow M$ (“universe rest frame”).
 - Pullback of metric and matter variables along \mathcal{X} .
 - Velocity of the source matter: ${}^{\mathcal{X}}v^i = {}^{\mathcal{X}}u^i / {}^{\mathcal{X}}u^0$.
 - Assume that source matter is slow-moving: $|{}^{\mathcal{X}}\vec{v}| \ll 1$.
- Use $\epsilon = |{}^{\mathcal{X}}\vec{v}|$ as perturbation parameter.
- Assign velocity orders $\mathcal{O}(n) \sim \epsilon^n$ to all quantities.
- Quasi-static: assign additional $\mathcal{O}(1)$ to time derivatives ∂_0 .

Standard post-Newtonian metric

- PPN formalism assumes fixed standard gauge \mathcal{P} .

Standard post-Newtonian metric

- PPN formalism assumes fixed standard gauge \mathcal{P} .
- Metric in standard PPN gauge:

$${}^{\mathcal{P}}g_{00}^2 = 2^{\mathcal{P}}U,$$

$${}^{\mathcal{P}}g_{ij}^2 = 2\gamma^{\mathcal{P}}U\delta_{ij},$$

$${}^{\mathcal{P}}g_{0i}^3 = -\frac{1}{2}(3 + 4\gamma + \alpha_1 - \alpha_2 + \zeta_1 - 2\xi)^{\mathcal{P}}V_i - \frac{1}{2}(1 + \alpha_2 - \zeta_1 + 2\xi)^{\mathcal{P}}W_i,$$

$$\begin{aligned} {}^{\mathcal{P}}g_{00}^4 &= (2 + 2\gamma + \alpha_3 + \zeta_1 - 2\xi)^{\mathcal{P}}\Phi_1 + 2(1 + 3\gamma - 2\beta + \zeta_2 + \xi)^{\mathcal{P}}\Phi_2 \\ &\quad + 2(1 + \zeta_3)^{\mathcal{P}}\Phi_3 + 2(3\gamma + 3\zeta_4 - 2\xi)^{\mathcal{P}}\Phi_4 - 2\xi^{\mathcal{P}}\Phi_W \\ &\quad - (\zeta_1 - 2\xi)^{\mathcal{P}}\mathfrak{A} - 2\beta^{\mathcal{P}}U^2. \end{aligned}$$

Standard post-Newtonian metric

- PPN formalism assumes fixed standard gauge \mathcal{P} .
- Metric in standard PPN gauge:

$${}^{\mathcal{P}}g_{00} = 2^{\mathcal{P}}U,$$

$${}^{\mathcal{P}}g_{ij} = 2\gamma^{\mathcal{P}}U\delta_{ij},$$

$${}^{\mathcal{P}}g_{0i} = -\frac{1}{2}(3 + 4\gamma + \alpha_1 - \alpha_2 + \zeta_1 - 2\xi)^{\mathcal{P}}V_i - \frac{1}{2}(1 + \alpha_2 - \zeta_1 + 2\xi)^{\mathcal{P}}W_i,$$

$$\begin{aligned} {}^{\mathcal{P}}g_{00} = & (2 + 2\gamma + \alpha_3 + \zeta_1 - 2\xi)^{\mathcal{P}}\Phi_1 + 2(1 + 3\gamma - 2\beta + \zeta_2 + \xi)^{\mathcal{P}}\Phi_2 \\ & + 2(1 + \zeta_3)^{\mathcal{P}}\Phi_3 + 2(3\gamma + 3\zeta_4 - 2\xi)^{\mathcal{P}}\Phi_4 - 2\xi^{\mathcal{P}}\Phi_W \\ & - (\zeta_1 - 2\xi)^{\mathcal{P}}\mathfrak{A} - 2\beta^{\mathcal{P}}U^2. \end{aligned}$$

- Metric contains **PPN parameters** and **PPN potentials**.
 - **PPN potentials** describe source matter distribution.
 - **PPN parameters** characterize gravity theory.

Standard post-Newtonian metric

- PPN formalism assumes fixed standard gauge \mathcal{P} .
- Metric in standard PPN gauge:

$${}^{\mathcal{P}}g_{00}^2 = 2^{\mathcal{P}}U,$$

$${}^{\mathcal{P}}g_{ij}^2 = 2\gamma^{\mathcal{P}}U\delta_{ij},$$

$${}^{\mathcal{P}}g_{0i}^3 = -\frac{1}{2}(3 + 4\gamma + \alpha_1 - \alpha_2 + \zeta_1 - 2\xi)^{\mathcal{P}}V_i - \frac{1}{2}(1 + \alpha_2 - \zeta_1 + 2\xi)^{\mathcal{P}}W_i,$$

$$\begin{aligned} {}^{\mathcal{P}}g_{00}^4 &= (2 + 2\gamma + \alpha_3 + \zeta_1 - 2\xi)^{\mathcal{P}}\Phi_1 + 2(1 + 3\gamma - 2\beta + \zeta_2 + \xi)^{\mathcal{P}}\Phi_2 \\ &\quad + 2(1 + \zeta_3)^{\mathcal{P}}\Phi_3 + 2(3\gamma + 3\zeta_4 - 2\xi)^{\mathcal{P}}\Phi_4 - 2\xi^{\mathcal{P}}\Phi_W \\ &\quad - (\zeta_1 - 2\xi)^{\mathcal{P}}\mathfrak{A} - 2\beta^{\mathcal{P}}U^2. \end{aligned}$$

- Metric contains PPN parameters and PPN potentials.
 - PPN potentials describe source matter distribution.
 - PPN parameters characterize gravity theory.
- ↪ Decompose metric into gauge-invariant and pure gauge parts.

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations
- 3 Parametrized post-Newtonian formalism
- 4 Gauge-invariant PPN formalism**
- 5 Example: PPN limit of scalar-tensor gravity
- 6 Conclusion

Gauge-invariant metric

- Definition of gauge-invariant metric components:

$$\mathbf{g}_{00} = \mathbf{g}^{\star}, \quad \mathbf{g}_{0i} = \mathbf{g}_i^{\diamond}, \quad \mathbf{g}_{ij} = \mathbf{g}^{\bullet} \delta_{ij} + \mathbf{g}_{ij}^{\dagger}.$$

Gauge-invariant metric

- Definition of gauge-invariant metric components:

$$\mathbf{g}_{00} = \mathbf{g}^{\star}, \quad \mathbf{g}_{0i} = \mathbf{g}_i^{\diamond}, \quad \mathbf{g}_{ij} = \mathbf{g}^{\bullet} \delta_{ij} + \mathbf{g}_{ij}^{\dagger}.$$

- Conditions imposed on components:

$$\partial^i \mathbf{g}_i^{\diamond} = 0, \quad \partial^i \mathbf{g}_{ij}^{\dagger} = 0, \quad \mathbf{g}_{[ij]}^{\dagger} = 0, \quad \mathbf{g}_{ii}^{\dagger} = 0.$$

Gauge-invariant metric

- Definition of gauge-invariant metric components:

$$\mathbf{g}_{00} = \mathbf{g}^*, \quad \mathbf{g}_{0i} = \mathbf{g}_i^\diamond, \quad \mathbf{g}_{ij} = \mathbf{g}^\bullet \delta_{ij} + \mathbf{g}_{ij}^\dagger.$$

- Conditions imposed on components:

$$\partial^i \mathbf{g}_i^\diamond = 0, \quad \partial^i \mathbf{g}_{ij}^\dagger = 0, \quad \mathbf{g}_{[ij]}^\dagger = 0, \quad \mathbf{g}_{ii}^\dagger = 0.$$

- Relation to arbitrary gauge \mathcal{X} :

$$\mathcal{X}^2 \mathbf{g}_{00} = \mathbf{g}^{2*},$$

$$\mathcal{X}^2 \mathbf{g}_{ij} = \mathbf{g}^{2\bullet} \delta_{ij} + \mathbf{g}_{ij}^{2\dagger} + 2\partial_i \partial_j \mathcal{X}^{2\diamond} + 2\partial_{(i} \mathcal{X}_{j)}^{2\diamond},$$

$$\mathcal{X}^3 \mathbf{g}_{0i} = \mathbf{g}_i^{3\diamond} + \partial_i \mathcal{X}^{3*} + \partial_0 \partial_i \mathcal{X}^{2\diamond} + \partial_0 \mathcal{X}_i^{2\diamond},$$

$$\mathcal{X}^4 \mathbf{g}_{00} = \mathbf{g}^{4*} + 2\partial_0 \mathcal{X}^{3*} + (\partial_i \mathcal{X}^{2\diamond} + \mathcal{X}_i^{2\diamond}) \partial_i \mathbf{g}^{2*},$$

$$\mathcal{X}^4 \mathbf{g}_{ij} = \mathbf{g}^{4\bullet} \delta_{ij} + \mathbf{g}_{ij}^{4\dagger} + 2\partial_i \partial_j \mathcal{X}^{4\diamond} + \mathcal{O}(2) \cdot \mathcal{O}(2).$$

Gauge-invariant metric

- Definition of gauge-invariant metric components:

$$\mathbf{g}_{00} = \mathbf{g}^*, \quad \mathbf{g}_{0i} = \mathbf{g}_i^\diamond, \quad \mathbf{g}_{ij} = \mathbf{g}^\bullet \delta_{ij} + \mathbf{g}_{ij}^\dagger.$$

- Conditions imposed on components:

$$\partial^i \mathbf{g}_i^\diamond = 0, \quad \partial^i \mathbf{g}_{ij}^\dagger = 0, \quad \mathbf{g}_{[ij]}^\dagger = 0, \quad \mathbf{g}_{ii}^\dagger = 0.$$

- Relation to arbitrary gauge \mathcal{X} :

$$\mathcal{X}^2 \mathbf{g}_{00} = \mathbf{g}^{2*},$$

$$\mathcal{X}^2 \mathbf{g}_{ij} = \mathbf{g}^{2\bullet} \delta_{ij} + \mathbf{g}_{ij}^{2\dagger} + 2\partial_i \partial_j \mathcal{X}^{2\diamond} + 2\partial_{(i} \mathcal{X}_{j)}^{2\diamond},$$

$$\mathcal{X}^3 \mathbf{g}_{0i} = \mathbf{g}_i^{3\diamond} + \partial_i \mathcal{X}^{3*} + \partial_0 \partial_i \mathcal{X}^{2\diamond} + \partial_0 \mathcal{X}_i^{2\diamond},$$

$$\mathcal{X}^4 \mathbf{g}_{00} = \mathbf{g}^{4*} + 2\partial_0 \mathcal{X}^{3*} + (\partial_i \mathcal{X}^{2\diamond} + \mathcal{X}_i^{2\diamond}) \partial_i \mathbf{g}^{2*},$$

$$\mathcal{X}^4 \mathbf{g}_{ij} = \mathbf{g}^{4\bullet} \delta_{ij} + \mathbf{g}_{ij}^{4\dagger} + 2\partial_i \partial_j \mathcal{X}^{4\diamond} + \mathcal{O}(2) \cdot \mathcal{O}(2).$$

- Gauge defining vector fields:

$$X_i = \partial_i \mathcal{X}^\diamond + X_i^\diamond, \quad X_0 = \mathcal{X}^*, \quad \partial^i X_i^\diamond = 0.$$

Decomposition of metric components

- Count number of independent components at each order:

	total		invariant		pure gauge
$x^2 g_{00}$	1	\mathbf{g}^*	1	-	0
$x^2 g_{ij}$	6	$\mathbf{g}^*, \mathbf{g}_{ij}^\dagger$	1 + 2	X^\diamond, X_i^\diamond	1 + 2
$x^3 g_{0i}$	3	\mathbf{g}_i^\diamond	2	X^*	1
$x^4 g_{00}$	1	\mathbf{g}^*	1	-	0
$x^4 g_{ij}$	6	$\mathbf{g}^*, \mathbf{g}_{ij}^\dagger$	1 + 2	X^\diamond, X_i^\diamond	1 + 2

Decomposition of metric components

- Count number of independent components at each order:

	total	invariant		pure gauge	
$x^2 g_{00}$	1	g^*	1	-	0
$x^2 g_{ij}$	6	g^*, g_{ij}^\dagger	1 + 2	X^\diamond, X_i^\diamond	1 + 2
$x^3 g_{0i}$	3	g_i^\diamond	2	X^*	1
$x^4 g_{00}$	1	g^*	1	-	0
$x^4 g_{ij}$	6	g^*, g_{ij}^\dagger	1 + 2	X^\diamond, X_i^\diamond	1 + 2

⇒ Components split into invariant and gauge parts.

Decomposition of metric components

- Count number of independent components at each order:

	total	invariant		pure gauge	
$x^2 g_{00}$	1	g^*	1	-	0
$x^2 g_{ij}$	6	g^*, g_{ij}^\dagger	1 + 2	X^\diamond, X_i^\diamond	1 + 2
$x^3 g_{0i}$	3	g_i^\diamond	2	X^*	1
$x^4 g_{00}$	1	g^*	1	-	0
$x^4 g_{ij}$	6	g^*, g_{ij}^\dagger	1 + 2	X^\diamond, X_i^\diamond	1 + 2

- ⇒ Components split into invariant and gauge parts.
- ⇒ Possible to separate physical information from coordinate choice.

Relation to standard PPN gauge

- Use relation between expansion coefficients:

$${}^{\mathcal{P}}\mathbf{g}^k = \sum_{0 \leq l_1 + 2l_2 + \dots \leq k} \frac{1}{l_1! l_2! \dots} \xi_{\mathcal{P}}^{l_1} \dots \xi_{\mathcal{P}}^{l_k} \dots \mathbf{g}^{k-l_1-2l_2-\dots}$$

Relation to standard PPN gauge

- Use relation between expansion coefficients:

$${}^{\mathcal{P}}\mathbf{g}^k = \sum_{0 \leq l_1 + 2l_2 + \dots \leq k} \frac{1}{l_1! l_2! \dots} \xi_{\mathcal{P}}^{l_1} \dots \xi_{\mathcal{P}}^{l_k} \dots \mathbf{g}^{k-l_1-2l_2-\dots}$$

- Split components of ${}^{\mathcal{P}}g_{\mu\nu}$ into $\mathbf{g}_{\mu\nu}$ and P^μ .

Relation to standard PPN gauge

- Use relation between expansion coefficients:

$${}^{\mathcal{P}}\mathbf{g}^k = \sum_{0 \leq l_1 + 2l_2 + \dots \leq k} \frac{1}{l_1! l_2! \dots} \xi_{\mathcal{P}}^{l_1} \dots \xi_{\mathcal{P}}^{l_k} \dots \mathbf{g}^{k-l_1-2l_2-\dots}$$

- Split components of ${}^{\mathcal{P}}g_{\mu\nu}$ into $\mathbf{g}_{\mu\nu}$ and P^μ .
- ⇒ Gauge defining vector fields:

$${}^2P^\diamond = 0, \quad {}^2P_j^\diamond = 0, \quad {}^3P^\star = -\frac{1}{4}(2 + 4\gamma + \alpha_1 - 2\alpha_2 + 2\zeta_1 - 4\xi)\chi_{,0}.$$

Relation to standard PPN gauge

- Use relation between expansion coefficients:

$${}^{\mathcal{P}}\mathbf{g}^k = \sum_{0 \leq l_1 + 2l_2 + \dots \leq k} \frac{1}{l_1! l_2! \dots} \xi_1^{l_1} \dots \xi_k^{l_k} \dots {}^k \mathbf{g}.$$

- Split components of ${}^{\mathcal{P}}g_{\mu\nu}$ into $\mathbf{g}_{\mu\nu}$ and P^μ .

⇒ Gauge defining vector fields:

$${}^2 P^\diamond = 0, \quad {}^2 P_i^\diamond = 0, \quad {}^3 P^\star = -\frac{1}{4}(2 + 4\gamma + \alpha_1 - 2\alpha_2 + 2\zeta_1 - 4\xi)\chi_{,0}.$$

⇒ Gauge-invariant metric components:

$${}^2 \mathbf{g}^\star = 2\mathbf{U}, \quad {}^2 \mathbf{g}^\bullet = 2\gamma\mathbf{U}, \quad {}^2 \mathbf{g}_{ij}^\dagger = 0, \quad {}^3 \mathbf{g}_i^\diamond = -\left(1 + \gamma + \frac{\alpha_1}{4}\right)(\mathbf{V}_i + \mathbf{W}_i),$$

$$\begin{aligned} {}^4 \mathbf{g}^\star &= \frac{1}{2}(2 - \alpha_1 + 2\alpha_2 + 2\alpha_3)\Phi_1 + 2(1 + 3\gamma - 2\beta + \zeta_2 + \xi)\Phi_2 \\ &\quad + 2(1 + \zeta_3)\Phi_3 + 2(3\gamma + 3\zeta_4 - 2\xi)\Phi_4 - 2\xi\Phi_W - 2\beta\mathbf{U}^2 \\ &\quad + \frac{1}{2}(2 + 4\gamma + \alpha_1 - 2\alpha_2)\mathfrak{A} + \frac{1}{2}(2 + 4\gamma + \alpha_1 - 2\alpha_2 + 2\zeta_1 - 4\xi)\mathfrak{B}. \end{aligned}$$

Gauge-invariant field equations

- Perform similar decomposition of energy-momentum tensor:

$$\mathbf{T}^\star = \mathbf{T}_{00} = \rho \left(1 - \mathbf{g}_{00}^2 + \mathbf{v}^2 + \boldsymbol{\Pi} \right) + \mathcal{O}(6),$$

$$\mathbf{T}_i^\diamond + \partial_i \mathbf{T}^\blacklozenge = \mathbf{T}_{0i} = -\rho \mathbf{v}_i + \mathcal{O}(5),$$

$$\mathbf{T}^\bullet \delta_{ij} + \Delta_{ij} \mathbf{T}^\blacktriangle + 2\partial_{(i} \mathbf{T}_{j)}^\blacktriangle + \mathbf{T}_{ij}^\dagger = \mathbf{T}_{ij} = \rho \mathbf{v}_i \mathbf{v}_j + \mathbf{p} \delta_{ij} + \mathcal{O}(6).$$

Gauge-invariant field equations

- Perform similar decomposition of energy-momentum tensor:

$$\mathbf{T}^* = \mathbf{T}_{00} = \rho \left(1 - \overset{2}{\mathbf{g}}_{00} + \mathbf{v}^2 + \boldsymbol{\Pi} \right) + \mathcal{O}(6),$$

$$\mathbf{T}_i^\diamond + \partial_i \mathbf{T}^\blacklozenge = \mathbf{T}_{0i} = -\rho \mathbf{v}_i + \mathcal{O}(5),$$

$$\mathbf{T}^\bullet \delta_{ij} + \Delta_{ij} \mathbf{T}^\blacktriangle + 2\partial_{(i} \mathbf{T}_{j)}^\blacktriangle + \mathbf{T}_{ij}^\dagger = \mathbf{T}_{ij} = \rho \mathbf{v}_i \mathbf{v}_j + \mathbf{p} \delta_{ij} + \mathcal{O}(6).$$

- Express components in terms of PPN potentials:

$$\overset{2}{\mathbf{T}}^* = \rho = -\frac{1}{4\pi} \Delta \mathbf{U}, \quad \overset{3}{\mathbf{T}}^\blacklozenge = -\frac{1}{4\pi} \partial_0 \mathbf{U}, \quad \overset{3}{\mathbf{T}}_i^\diamond = \frac{1}{8\pi} \Delta (\mathbf{V}_i + \mathbf{W}_i),$$

$$\overset{4}{\mathbf{T}}^* = \rho \left(\boldsymbol{\Pi} + \mathbf{v}^2 - \overset{2}{\mathbf{g}}^* \right) = -\frac{1}{4\pi} \Delta (\boldsymbol{\Phi}_3 + \boldsymbol{\Phi}_1 - 2\boldsymbol{\Phi}_2),$$

$$\overset{4}{\mathbf{T}}^\bullet = \frac{1}{3} \rho \mathbf{v}^2 + \mathbf{p} = -\frac{1}{12\pi} \Delta (\boldsymbol{\Phi}_1 + 3\boldsymbol{\Phi}_4), \quad \overset{4}{\mathbf{T}}^\blacktriangle = \frac{1}{16\pi} (3\boldsymbol{\alpha} - \boldsymbol{\Phi}_1).$$

Gauge-invariant field equations

- Perform similar decomposition of energy-momentum tensor:

$$\mathbf{T}^* = \mathbf{T}_{00} = \rho \left(1 - \overset{2}{\mathbf{g}}_{00} + \mathbf{v}^2 + \boldsymbol{\Pi} \right) + \mathcal{O}(6),$$

$$\mathbf{T}_i^\diamond + \partial_i \mathbf{T}^\blacklozenge = \mathbf{T}_{0i} = -\rho \mathbf{v}_i + \mathcal{O}(5),$$

$$\mathbf{T}^\bullet \delta_{ij} + \Delta_{ij} \mathbf{T}^\blacktriangle + 2\partial_{(i} \mathbf{T}_{j)}^\blacktriangle + \mathbf{T}_{ij}^\dagger = \mathbf{T}_{ij} = \rho \mathbf{v}_i \mathbf{v}_j + \mathbf{p} \delta_{ij} + \mathcal{O}(6).$$

- Express components in terms of PPN potentials:

$$\overset{2}{\mathbf{T}}^* = \rho = -\frac{1}{4\pi} \Delta \mathbf{U}, \quad \overset{3}{\mathbf{T}}^\blacklozenge = -\frac{1}{4\pi} \partial_0 \mathbf{U}, \quad \overset{3}{\mathbf{T}}_i^\diamond = \frac{1}{8\pi} \Delta (\mathbf{V}_i + \mathbf{W}_i),$$

$$\overset{4}{\mathbf{T}}^* = \rho \left(\boldsymbol{\Pi} + \mathbf{v}^2 - \overset{2}{\mathbf{g}}^* \right) = -\frac{1}{4\pi} \Delta (\boldsymbol{\Phi}_3 + \boldsymbol{\Phi}_1 - 2\boldsymbol{\Phi}_2),$$

$$\overset{4}{\mathbf{T}}^\bullet = \frac{1}{3} \rho \mathbf{v}^2 + \mathbf{p} = -\frac{1}{12\pi} \Delta (\boldsymbol{\Phi}_1 + 3\boldsymbol{\Phi}_4), \quad \overset{4}{\mathbf{T}}^\blacktriangle = \frac{1}{16\pi} (3\boldsymbol{\alpha} - \boldsymbol{\Phi}_1).$$

- Decompose also gravity side $\simeq \Delta \mathbf{g}$.

Gauge-invariant field equations

- Perform similar decomposition of energy-momentum tensor:

$$\mathbf{T}^* = \mathbf{T}_{00} = \rho \left(1 - \overset{2}{\mathbf{g}}_{00} + \mathbf{v}^2 + \boldsymbol{\Pi} \right) + \mathcal{O}(6),$$

$$\mathbf{T}_i^\diamond + \partial_i \mathbf{T}^\blacklozenge = \mathbf{T}_{0i} = -\rho \mathbf{v}_i + \mathcal{O}(5),$$

$$\mathbf{T}^\bullet \delta_{ij} + \Delta_{ij} \mathbf{T}^\blacktriangle + 2\partial_{(i} \mathbf{T}_{j)}^\blacktriangle + \mathbf{T}_{ij}^\dagger = \mathbf{T}_{ij} = \rho \mathbf{v}_i \mathbf{v}_j + \mathbf{p} \delta_{ij} + \mathcal{O}(6).$$

- Express components in terms of PPN potentials:

$$\overset{2}{\mathbf{T}}^* = \rho = -\frac{1}{4\pi} \Delta \mathbf{U}, \quad \overset{3}{\mathbf{T}}^\blacklozenge = -\frac{1}{4\pi} \partial_0 \mathbf{U}, \quad \overset{3}{\mathbf{T}}_i^\diamond = \frac{1}{8\pi} \Delta (\mathbf{V}_i + \mathbf{W}_i),$$

$$\overset{4}{\mathbf{T}}^* = \rho \left(\boldsymbol{\Pi} + \mathbf{v}^2 - \overset{2}{\mathbf{g}}^* \right) = -\frac{1}{4\pi} \Delta (\boldsymbol{\Phi}_3 + \boldsymbol{\Phi}_1 - 2\boldsymbol{\Phi}_2),$$

$$\overset{4}{\mathbf{T}}^\bullet = \frac{1}{3} \rho \mathbf{v}^2 + \mathbf{p} = -\frac{1}{12\pi} \Delta (\boldsymbol{\Phi}_1 + 3\boldsymbol{\Phi}_4), \quad \overset{4}{\mathbf{T}}^\blacktriangle = \frac{1}{16\pi} (3\boldsymbol{\alpha} - \boldsymbol{\Phi}_1).$$

- Decompose also gravity side $\simeq \Delta \mathbf{g}$.

⇒ Find PPN parameters by comparing coefficients on both sides.

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations
- 3 Parametrized post-Newtonian formalism
- 4 Gauge-invariant PPN formalism
- 5 Example: PPN limit of scalar-tensor gravity**
- 6 Conclusion

Action and field equations

- Action of scalar-tensor gravity with massless scalar field: [Nordtvedt '70]

$$S = \frac{1}{2\kappa^2} \int_M d^4x \sqrt{-g} \left(\psi R - \frac{\omega(\psi)}{\psi} \partial_\rho \psi \partial^\rho \psi \right) + S_m[g_{\mu\nu}, \chi].$$

Action and field equations

- Action of scalar-tensor gravity with massless scalar field: [Nordtvedt '70]

$$S = \frac{1}{2\kappa^2} \int_M d^4x \sqrt{-g} \left(\psi R - \frac{\omega(\psi)}{\psi} \partial_\rho \psi \partial^\rho \psi \right) + S_m[g_{\mu\nu}, \chi].$$

- Free function $\omega(\psi)$ of the scalar field ψ .

Action and field equations

- Action of scalar-tensor gravity with massless scalar field: [Nordtvedt '70]

$$S = \frac{1}{2\kappa^2} \int_M d^4x \sqrt{-g} \left(\psi R - \frac{\omega(\psi)}{\psi} \partial_\rho \psi \partial^\rho \psi \right) + S_m[g_{\mu\nu}, \chi].$$

- Free function $\omega(\psi)$ of the scalar field ψ .
- Jordan frame: no coupling between matter and scalar field.

Action and field equations

- Action of scalar-tensor gravity with massless scalar field: [Nordtvedt '70]

$$S = \frac{1}{2\kappa^2} \int_M d^4x \sqrt{-g} \left(\psi R - \frac{\omega(\psi)}{\psi} \partial_\rho \psi \partial^\rho \psi \right) + S_m[g_{\mu\nu}, \chi].$$

- Free function $\omega(\psi)$ of the scalar field ψ .
- Jordan frame: no coupling between matter and scalar field.

⇒ Field equations:

$$\begin{aligned} \psi R_{\mu\nu} - \nabla_\mu \partial_\nu \psi - \frac{\omega}{\psi} \partial_\mu \psi \partial_\nu \psi + \frac{g_{\mu\nu}}{4\omega + 6} \frac{d\omega}{d\psi} \partial_\rho \psi \partial^\rho \psi \\ = \kappa^2 \left(T_{\mu\nu} - \frac{\omega + 1}{2\omega + 3} g_{\mu\nu} T \right), \end{aligned}$$

$$(2\omega + 3) \square \psi + \frac{d\omega}{d\psi} \partial_\rho \psi \partial^\rho \psi = \kappa^2 T.$$

Action and field equations

- Action of scalar-tensor gravity with massless scalar field: [Nordtvedt '70]

$$S = \frac{1}{2\kappa^2} \int_M d^4x \sqrt{-g} \left(\psi R - \frac{\omega(\psi)}{\psi} \partial_\rho \psi \partial^\rho \psi \right) + S_m[g_{\mu\nu}, \chi].$$

- Free function $\omega(\psi)$ of the scalar field ψ .
- Jordan frame: no coupling between matter and scalar field.

⇒ Field equations:

$$\begin{aligned} \psi R_{\mu\nu} - \nabla_\mu \partial_\nu \psi - \frac{\omega}{\psi} \partial_\mu \psi \partial_\nu \psi + \frac{g_{\mu\nu}}{4\omega + 6} \frac{d\omega}{d\psi} \partial_\rho \psi \partial^\rho \psi \\ = \kappa^2 \left(T_{\mu\nu} - \frac{\omega + 1}{2\omega + 3} g_{\mu\nu} T \right), \end{aligned}$$

$$(2\omega + 3) \square \psi + \frac{d\omega}{d\psi} \partial_\rho \psi \partial^\rho \psi = \kappa^2 T.$$

↪ Decompose into gauge-invariant field equations.

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .
- Relevant components of scalar field: $x^0_\psi = \Psi, x^2_\psi, x^4_\psi$.

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .
- Relevant components of scalar field: $x^0_\psi = \Psi$, x^2_ψ , x^4_ψ .
- Cosmological background value Ψ assumed to be constant.

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .
- Relevant components of scalar field: $x^0_\psi = \Psi$, x^2_ψ , x^4_ψ .
- Cosmological background value Ψ assumed to be constant.
- Relation to gauge-invariant scalar field perturbations $\overset{2}{\psi}$ and $\overset{2}{\psi}$:

$$x^2_\psi = \overset{2}{\psi}, \quad x^4_\psi = \overset{4}{\psi} + (\partial_i \overset{2}{X}^\diamond + \overset{2}{X}^\diamond_{,i}) \overset{2}{\psi}_{,i}.$$

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .
- Relevant components of scalar field: $x^0_\psi = \Psi$, x^2_ψ , x^4_ψ .
- Cosmological background value Ψ assumed to be constant.
- Relation to gauge-invariant scalar field perturbations $\overset{2}{\psi}$ and $\overset{2}{\psi}$:

$$x^2_\psi = \overset{2}{\psi}, \quad x^4_\psi = \overset{4}{\psi} + (\partial_i \overset{2}{X}^\diamond + \overset{2}{X}^\diamond_i) \overset{2}{\psi}_{,i}.$$

- Taylor expansion of function ω around background value:

$$\omega_0 = \omega(\Psi), \quad \omega_1 = \omega'(\Psi).$$

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .
- Relevant components of scalar field: $\mathcal{X}^0 \psi = \Psi$, $\mathcal{X}^2 \psi$, $\mathcal{X}^4 \psi$.
- Cosmological background value Ψ assumed to be constant.
- Relation to gauge-invariant scalar field perturbations $\overset{2}{\psi}$ and $\overset{2}{\psi}$:

$$\mathcal{X}^2 \psi = \overset{2}{\psi}, \quad \mathcal{X}^4 \psi = \overset{4}{\psi} + (\partial_i \overset{2}{X}^\diamond + \overset{2}{X}_i^\diamond) \overset{2}{\psi}_{,i}.$$

- Taylor expansion of function ω around background value:

$$\omega_0 = \omega(\Psi), \quad \omega_1 = \omega'(\Psi).$$

⇒ Zeroth order $\mathcal{X}^0 \psi = \Psi$, $\mathcal{X}^0 \overset{0}{g}_{\mu\nu} = \eta_{\mu\nu}$ solves (vacuum) field equations.

Perturbative solution ansatz

- Consider gauge \mathcal{X} to pull equations and fields to M_0 .
- Relevant components of scalar field: $\mathcal{X}^0 \psi = \Psi$, $\mathcal{X}^2 \psi$, $\mathcal{X}^4 \psi$.
- Cosmological background value Ψ assumed to be constant.
- Relation to gauge-invariant scalar field perturbations $\overset{2}{\psi}$ and $\overset{2}{\psi}$:

$$\mathcal{X}^2 \psi = \overset{2}{\psi}, \quad \mathcal{X}^4 \psi = \overset{4}{\psi} + (\partial_i \overset{2}{X}^\diamond + \overset{2}{X}_i^\diamond) \overset{2}{\psi}_{,i}.$$

- Taylor expansion of function ω around background value:

$$\omega_0 = \omega(\Psi), \quad \omega_1 = \omega'(\Psi).$$

- ⇒ Zeroth order $\mathcal{X}^0 \psi = \Psi$, $\mathcal{X}^0 \overset{0}{g}_{\mu\nu} = \eta_{\mu\nu}$ solves (vacuum) field equations.
- ⇒ Remaining equations determine gauge-invariant metric.

Post-Newtonian metric and PPN parameters

- Normalization of the gravitational constant:

$$\kappa^2 = 4\pi\Psi \frac{2\omega_0 + 3}{\omega_0 + 2} .$$

Post-Newtonian metric and PPN parameters

- Normalization of the gravitational constant:

$$\kappa^2 = 4\pi\Psi \frac{2\omega_0 + 3}{\omega_0 + 2}.$$

⇒ Metric in terms of PPN potentials:

$$\mathbf{g}^2_{\star} = 2\mathbf{U}, \quad \mathbf{g}^2_{\bullet} = 2\frac{\omega_0 + 1}{\omega_0 + 2}\mathbf{U}, \quad \mathbf{g}^2_{ij} = 0, \quad \mathbf{g}^3_{i\diamond} = -\frac{2\omega_0 + 3}{\omega_0 + 2}(\mathbf{V}_i + \mathbf{W}_i),$$

$$\mathbf{g}^4_{\star} = \frac{3\omega_0 + 4}{\omega_0 + 2}(\mathfrak{A} + \mathfrak{B}) + \Phi_1 + \left(\frac{4\omega_0 + 2}{\omega_0 + 2} - \frac{\omega_1\Psi}{(2\omega_0 + 3)(\omega_0 + 2)^2} \right) \Phi_2 \\ + 3\Phi_3 + 6\frac{\omega_0 + 1}{\omega_0 + 2}\Phi_4 - 2\left(1 + \frac{\omega_1\Psi}{4(2\omega_0 + 3)(\omega_0 + 2)^2} \right) \mathbf{U}^2.$$

Post-Newtonian metric and PPN parameters

- Normalization of the gravitational constant:

$$\kappa^2 = 4\pi\Psi \frac{2\omega_0 + 3}{\omega_0 + 2}.$$

- ⇒ Metric in terms of PPN potentials:

$$\mathbf{g}^2_* = 2\mathbf{U}, \quad \mathbf{g}^2_\bullet = 2\frac{\omega_0 + 1}{\omega_0 + 2}\mathbf{U}, \quad \mathbf{g}^2_{ij} = 0, \quad \mathbf{g}^3_i{}^\diamond = -\frac{2\omega_0 + 3}{\omega_0 + 2}(\mathbf{V}_i + \mathbf{W}_i),$$

$$\mathbf{g}^4_* = \frac{3\omega_0 + 4}{\omega_0 + 2}(\mathfrak{A} + \mathfrak{B}) + \Phi_1 + \left(\frac{4\omega_0 + 2}{\omega_0 + 2} - \frac{\omega_1\Psi}{(2\omega_0 + 3)(\omega_0 + 2)^2} \right) \Phi_2 \\ + 3\Phi_3 + 6\frac{\omega_0 + 1}{\omega_0 + 2}\Phi_4 - 2\left(1 + \frac{\omega_1\Psi}{4(2\omega_0 + 3)(\omega_0 + 2)^2} \right) \mathbf{U}^2.$$

- ⇒ PPN parameters reproduce well-known result: [Nordtvedt '70]

$$\gamma = \frac{\omega_0 + 1}{\omega_0 + 2}, \quad \beta = 1 + \frac{\omega_1\Psi}{4(2\omega_0 + 3)(\omega_0 + 2)^2},$$

$$\alpha_1 = \alpha_2 = \alpha_3 = \zeta_1 = \zeta_2 = \zeta_3 = \zeta_4 = \xi = 0.$$

Outline

- 1 Introduction
- 2 Gauge-invariant higher order perturbations
- 3 Parametrized post-Newtonian formalism
- 4 Gauge-invariant PPN formalism
- 5 Example: PPN limit of scalar-tensor gravity
- 6 Conclusion**

Summary

- Gauge-invariant perturbation theory:
 - Distinguish between physical and background spacetime.
 - Gauge pulls physical metric to background spacetime.
 - ⇒ Gauge dependent comparison between both metrics.
 - Decompose perturbations into physical data and gauge data.

Summary

- Gauge-invariant perturbation theory:
 - Distinguish between physical and background spacetime.
 - Gauge pulls physical metric to background spacetime.
 - ⇒ Gauge dependent comparison between both metrics.
 - Decompose perturbations into physical data and gauge data.
- Parametrized post-Newtonian formalism:
 - Weak-field approximation of metric gravity theories.
 - Characterizes gravity theories by 10 (constant) parameters.
 - Parameters closely related to solar system observations.

Summary

- Gauge-invariant perturbation theory:
 - Distinguish between physical and background spacetime.
 - Gauge pulls physical metric to background spacetime.
 - ⇒ Gauge dependent comparison between both metrics.
 - Decompose perturbations into physical data and gauge data.
- Parametrized post-Newtonian formalism:
 - Weak-field approximation of metric gravity theories.
 - Characterizes gravity theories by 10 (constant) parameters.
 - Parameters closely related to solar system observations.
- Gauge-invariant PPN formalism:
 - Apply gauge-invariant perturbation theory to PPN formalism.
 - Decomposition of metric and field equations.
 - Avoids issues arising from necessity to choose a gauge.
 - ⇒ Simpler set of equations to determine PPN parameters.
 - Can also be formulated for tetrad instead of metric.

Summary

- Gauge-invariant perturbation theory:
 - Distinguish between physical and background spacetime.
 - Gauge pulls physical metric to background spacetime.
 - ⇒ Gauge dependent comparison between both metrics.
 - Decompose perturbations into physical data and gauge data.
- Parametrized post-Newtonian formalism:
 - Weak-field approximation of metric gravity theories.
 - Characterizes gravity theories by 10 (constant) parameters.
 - Parameters closely related to solar system observations.
- Gauge-invariant PPN formalism:
 - Apply gauge-invariant perturbation theory to PPN formalism.
 - Decomposition of metric and field equations.
 - Avoids issues arising from necessity to choose a gauge.
 - ⇒ Simpler set of equations to determine PPN parameters.
 - Can also be formulated for tetrad instead of metric.
- Post-Newtonian limit of scalar-tensor gravity:
 - Perturbative field equations simplify in gauge-invariant formulation.
 - Consistency check: obtain well-known PPN parameters.
 - Also possible to use tetrad formulation to calculate solution.

- Extend formalism by including higher perturbation orders:
 - General covariant expansion instead of space-time split.
 - Allow also for fast-moving source masses.
 - Consider inspiral phase of black hole merger event.
 - Devise method for calculating gravitational waves.

- Extend formalism by including higher perturbation orders:
 - General covariant expansion instead of space-time split.
 - Allow also for fast-moving source masses.
 - Consider inspiral phase of black hole merger event.
 - Devise method for calculating gravitational waves.
- Further possible extensions and modifications:
 - Apply to formalism based on modified density $\rho^* = \rho\sqrt{-g}u^0$. [Will '18]
 - Allow for multiple dynamical metrics / tetrads.
 - Take into account Vainshtein or other screening mechanisms.
 - Cosmological background and time variability of PPN parameters.
 - Include PPN potentials appearing in higher derivative theories.

- Extend formalism by including higher perturbation orders:
 - General covariant expansion instead of space-time split.
 - Allow also for fast-moving source masses.
 - Consider inspiral phase of black hole merger event.
 - Devise method for calculating gravitational waves.
- Further possible extensions and modifications:
 - Apply to formalism based on modified density $\rho^* = \rho\sqrt{-g}u^0$. [Will '18]
 - Allow for multiple dynamical metrics / tetrads.
 - Take into account Vainshtein or other screening mechanisms.
 - Cosmological background and time variability of PPN parameters.
 - Include PPN potentials appearing in higher derivative theories.
- Apply formalism to complicated gravity theories:
 - Bimetric and multimetric gravity theories.
 - Multi-scalar Horndeski generalizations.
 - Theories involving generalized Proca fields.
 - Extensions based on metric-affine geometry.
 - Extensions of teleparallel and symmetric teleparallel gravity.

Further reading

MH,
“Gauge invariant approach to the parametrized post-Newtonian formalism”,
arXiv:1910.09245 [gr-qc] (to appear in Phys. Rev. D).